
Heavy-flavor measurements with the PHENIX experiment at RHIC

Ralf Averbeck

State University of New York at Stony Brook

for the **PHENIX** Collaboration

DPG Spring Meeting

Cologne, Germany, March 8-12, 2004

Outline

- Motivation
- PHENIX @ RHIC
- Open heavy-flavor (charm) measurements
 - method
 - selected results
- Heavy quarkonia (J/ψ) measurements
 - method
 - selected results
- Summary
- Outlook

Motivation I: open charm production

- charm yield in heavy-ion collisions

- production mainly via gluon-gluon fusion in the earliest stage of the collision



sensitive to initial gluon density

- additional thermal production at very high temperature
→ enhancement?



sensitive to initial temperature

- propagation through dense medium

- energy loss by gluon radiation?
→ softening of “charmed hadron” spectra?



sensitive to state of nuclear medium

- does charm flow?



sensitive to collectivity on partonic level




Motivation II: charmonia (J/ψ)

- $c\bar{c}$: produced early / embedded in medium
- can form bound state: J/ψ
- deconfinement & color screening
 $\Rightarrow J/\psi$ suppression (Matsui and Satz, PLB176(1986)416)
- central Pb+Pb collisions at SPS

- J/ψ suppression in excess of “normal”
nuclear suppression (NA50: PLB477(2000)28)


- prospects at RHIC

- higher $c\bar{c}$ yield than at SPS
- possible J/ψ enhancement due to $c\bar{c}$ coalescence as the medium cools
- important to measure J/ψ in p+p and d+A to separate “normal” nuclear effects
 - (anti) shadowing
 - nuclear absorption in cold matter
- open charm needed as baseline



PHENIX @ RHIC

- only RHIC experiment optimized for lepton measurements
- electrons: central arms
 - electron measurement in range:
 $|\eta| \leq 0.35$
 $p \geq 0.2 \text{ GeV}/c$
- muons: forward arms
 - muon measurement in range:
 $1.2 < |\eta| < 2.4$
 $p \geq 2 \text{ GeV}/c$





two central electron/photon/hadron spectrometers

two forward muon spectrometers

How to measure open charm


- ideal but very challenging

- direct reconstruction of charm decays (e.g. $D^0 \rightarrow K^- \pi^+$)



Open charm via single electrons (I)

- cocktail method (example)
 - inclusive e^\pm spectra from Au+Au at 130 GeV
 - use available data to establish “cocktail” of e^\pm sources
 - dominated by measured π^0 decays and photon conversions
 - “photonic sources” are photon conversions and Dalitz decays
- excess above cocktail
 - increasing with p_T
 - expected from charm decays
- subtract cocktail from data
- attribute excess to semileptonic decays of open charm
- first “measurement” of open charm in HI collisions



PHENIX: PRL 88(2002)192303


Open charm via single electrons (II)

- converter method

- add converter of known thickness to experiment
- compare e^\pm spectra with and without converter
- separation of e^\pm from photonic and non-photonic sources

- $e\gamma$ coincidence method



- e^\pm from photonic sources are correlated with a γ
- separation of e^\pm from different sources



- three independent methods to measure electrons from non-photonic sources!

Open charm in p+p collisions at $\sqrt{s} = 200$ GeV

- non-photonic e^\pm from p+p collisions at 200 GeV
- PYTHIA tuned to lower energy data ($\sqrt{s} < 63$ GeV)
- for $p_T > 1.5$ GeV/c spectra are “harder” than PYTHIA prediction



• reference for nuclear collisions

• spectral shape


- PYTHIA charm & bottom line shapes with normalizations as independent free parameters

• total cross section

- PYTHIA describes data at low p_T
- use PYTHIA to extrapolate to full phase space

Open charm from d+Au collisions at $\sqrt{s_{NN}} = 200$ GeV





PHENIX PRELIMINARY




- **non-photonic e^\pm from d+Au collisions at 200 GeV**

- **cross section divided by the nuclear overlap integral T_{AB} to account for the difference in system size between p+p and d+Au**
- **very good agreement of d+Au data with the scaled p+p reference within uncertainties**
- **no indication for strong cold-nuclear matter modifications**
- **how about centrality dependence?**

Centrality (in)dependence in d+Au collisions








Open charm from Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV



- **non-photonic e^\pm from Au+Au collisions at 200 GeV**


- **cross section divided by the nuclear overlap integral TAB to account for the difference in system size between p+p and Au+Au**
- **very good agreement of Au+Au data with the scaled p+p reference within uncertainties up to $p_T = 1.5$ GeV/c**
- **at higher p_T more statistics is needed**
- **no indication for strong medium modifications**
- **how about centrality dependence?**

Centrality (in)dependence in Au+Au collisions



● more statistics is highly desirable!


Binary collision scaling in Au+Au



- binary scaling seems to work!

- charm production seems to scale with T_{AB} , i.e. with the number of binary collisions N_{Coll}
- determine dN/dy of electrons in the measured range of p_T and test its consistency with $dN/dy = A (N_{\text{coll}})^\alpha$
- the 90 % Confidence Level on α is shown as yellow band

Does charm flow in Au+Au?



- uncertainties are too large to make a definite statement!



- charm flow could indicate partonic flow, i.e. collectivity on the parton level
- v_2 measures the 2nd harmonic of the Fourier expansion of the azimuthal distribution of particle emission with respect to the reaction plane
- v_2 of e^\pm from non-photonic sources is related to charm v_2 !

Summary (I): open charm at RHIC


- PHENIX: inclusive electrons in p+p, d+Au, and Au+Au at $\sqrt{s_{NN}} = 200$ GeV
- yield of electrons from non-photon sources
 - consistent with binary scaling
 - no indication for strong enhancement / suppression of charm cross section in nuclear collisions
- statistics limited regarding
 - presence of spectral modifications in Au+Au (energy loss)?
 - charm flow in Au+Au?
 - to be answered by Run-4 data (currently ongoing)
- charm data as baseline for J/ψ measurements!

How to identify J/ψ (example: d+Au @ 200 GeV)

- $J/\psi \rightarrow e^+e^-$



- $J/\psi \rightarrow \mu^+\mu^-$ (north μ -arm)



- subtraction of combinatorial background by subtraction of like-sign dilepton mass spectra from unlike-sign spectra
- (small) physical background remains

J/ψ suppression/enhancement in Au+Au at RHIC?

- preparing the case (need integrated luminosity!)

Year	Ions	$\sqrt{s_{NN}}$	Detectors	J/ψ
2000	Au-Au	130 GeV	Central (electrons)	0
2001	Au-Au	200 GeV	Central	13 + 0 [1]
2002	p-p	200 GeV	+ 1 muon arm	46 + 66 [2]
2002	d-Au	200 GeV	Central	300+800+600
2003	p-p	200 GeV	+ 2 muon arms	100+300+120
2004	Au-Au	200 GeV	! taking data !	~400+2x1600 ?

} first J/ψ
observation
at RHIC

} first sizeable
samples in
p+p & d+Au




- study modification of J/ψ production in cold nuclear matter: p+p versus d+Au!



$J/\psi \rightarrow \mu^+\mu^-$ Analysis for d+Au & p+p

- J/ψ mostly produced by gluon fusion
→ sensitive to gluon pdf
- three rapidity ranges probe different momentum fractions x_2 of Au partons
 - South ($y < -1.2$) : large x_2 (in gold) ~ 0.090
 - Central ($y \sim 0$) : intermediate x_2 ~ 0.020
 - North ($y > 1.2$) : small x_2 (in gold) ~ 0.003
- (anti)shadowing \leftrightarrow (enhancement) depletion of (high) low momentum partons

Example of predicted “gluon shadowing” in d+Au




From Eskola, Kolhinen, Vogt: Nucl. Phys. A696 (2001) 729-746.



J/ ψ ratio (d+Au / p+p) versus p_T


$$\sigma_{dA} = \sigma_{pp} (2 \times 197)^\alpha$$

- increase of α with increasing p_T
→ p_T broadening of J/ ψ (initial state multiple scattering, Cronin effect)
- similar to measurements at lower energy (E866: $\sqrt{s} = 39$ GeV)




J/ψ cross section versus rapidity

pp J/ψ – PHENIX Preliminary




dAu J/ψ – PHENIX Preliminary




$$BR \sigma_{pp}^{J/\psi} = 160 \text{ nb} \pm 8.5 \% (\text{fit}) \pm 12.3\% (\text{abs}) - \text{preliminary}$$

J/ ψ ratio (d+Au / p+p) versus rapidity

- data indicate nuclear effects (difficult to disentangle)!




• weak shadowing



• weak absorption

(central / peripheral) versus N_{coll}

$J/\Psi \rightarrow l^+l^-$ PHENIX Preliminary 200 GeV
Central/Peripheral (R_{cp}) vrs Number of Collisions



- define four centrality classes for d+Au collisions
- determine N_{coll} for each class in a Glauber model
- calculate R_{cp}

$$R_{\text{cp}}(N_{\text{coll}}) = \frac{N_{J\psi}^{\text{cent}} \times \langle N_{\text{coll}}^{\text{periph}} \rangle}{N_{J\psi}^{\text{periph}} \times \langle N_{\text{coll}}^{\text{cent}} \rangle}$$



- low and medium x_2
 - weak nuclear effects
 - small (shadowing) centrality dep.
- high x_2
 - steep rise with centrality
 - how can antishadowing rise so steeply while shadowing does not?
 - final state effect in Au remnants (close to Au frame)?

Summary (II): J/ψ at RHIC

- cold nuclear matter effects have been observed in d+Au collisions
 - weak gluon shadowing
 - weak absorption
 - p_T broadening: similar to lower energy data
- disentangling these effects requires more data
- modest baseline for J/ψ measurements in Au+Au collisions is available: the stage is set!
- Run-4 (Au+Au at 200 GeV) is currently ongoing!

Outlook

- J/ψ production in Au+Au at 200 GeV (Run-4)
- a first J/ψ signal in the dielectron channel
- stay tuned!



The PHENIX Collaboration

Brazil	University of São Paulo, São Paulo
China	Academia Sinica, Taipei, Taiwan China Institute of Atomic Energy, Beijing Peking University, Beijing
France	LPC, University de Clermont-Ferrand, Clermont-Ferrand Dapnia, CEA Saclay, Gif-sur-Yvette IPN-Orsay, Université Paris Sud, CNRS-IN2P3, Orsay LLR, École Polytechnique, CNRS-IN2P3, Palaiseau SUBATECH, École des Mines at Nantes, Nantes
Germany	University of Münster, Münster
Hungary	Central Research Institute for Physics (KFKI), Budapest Debrecen University, Debrecen Eötvös Loránd University (ELTE), Budapest
India	Banaras Hindu University, Banaras Bhabha Atomic Research Centre, Bombay
Israel	Weizmann Institute, Rehovot
Japan	Center for Nuclear Study, University of Tokyo, Tokyo Hiroshima University, Higashi-Hiroshima KEK, Institute for High Energy Physics, Tsukuba Kyoto University, Kyoto Nagasaki Institute of Applied Science, Nagasaki RIKEN, Institute for Physical and Chemical Research, Wako RIKEN-BNL Research Center, Upton, NY Rikkyo University, Tokyo, Japan Tokyo Institute of Technology, Tokyo University of Tsukuba, Tsukuba Waseda University, Tokyo
S. Korea	Cyclotron Application Laboratory, KAERI, Seoul Kangnung National University, Kangnung Korea University, Seoul Myong Ji University, Yongin City System Electronics Laboratory, Seoul Nat. University, Seoul Yonsei University, Seoul
Russia	Institute of High Energy Physics, Protovino Joint Institute for Nuclear Research, Dubna Kurchatov Institute, Moscow PNPI, St. Petersburg Nuclear Physics Institute, St. Petersburg St. Petersburg State Technical University, St. Petersburg
Sweden	Lund University, Lund



12 Countries; 58 Institutions; 480 Participants*

USA Abilene Christian University, Abilene, TX
Brookhaven National Laboratory, Upton, NY
University of California - Riverside, Riverside, CA
University of Colorado, Boulder, CO
Columbia University, Nevis Laboratories, Irvington, NY
Florida State University, Tallahassee, FL
Florida Technical University, Melbourne, FL
Georgia State University, Atlanta, GA
University of Illinois Urbana Champaign, Urbana-Champaign, IL
Iowa State University and Ames Laboratory, Ames, IA
Los Alamos National Laboratory, Los Alamos, NM
Lawrence Livermore National Laboratory, Livermore, CA
University of New Mexico, Albuquerque, NM
New Mexico State University, Las Cruces, NM
Dept. of Chemistry, Stony Brook Univ., Stony Brook, NY
Dept. Phys. and Astronomy, Stony Brook Univ., Stony Brook, NY
Oak Ridge National Laboratory, Oak Ridge, TN
University of Tennessee, Knoxville, TN
Vanderbilt University, Nashville, TN

***as of January 2004**